



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ALGEBRA.

Conducted by J. M. COLAW, Monterey, Va. All contributions to this department should be sent to him.

SOLUTIONS OF PROBLEMS.

60. Proposed by ROBERT J. ALEY, A. M., Professor of Mathematics in Indiana University, Bloomington, Indiana.

Telegraph poles are a yards apart; for how many minutes must one count poles in order that the number of poles counted may be equal to the number of miles per hour that the train is running?

I. Solution by FREDERICK R. HONEY, A. B., New Haven, Connecticut.

The problem is independent of the number of poles counted, and of the number of miles per hour the train is running. We will call this number N .

$\therefore aN$ = the number of yards the train runs while the poles are counted. Also, $1760N$ = number of yards per hour the train runs. $\therefore aN/1760N$ = the fraction of an hour during which the poles are counted.

$\therefore 60aN/1760N = 3a/88$ = number of minutes during which the poles are counted.

II. Solution by M. A. GRUBER, A. M., War Department, Washington, D. C.; and W. H. CARTER, Professor of Mathematics, Centenary College of Louisiana, Jackson, Louisiana.

Let x = the number of minutes, and let r = number of miles per hour the train is running. Also, $1760/a$ = number of poles in a mile, and $rx/60$ = number of miles the train runs in x minutes. Then, $rx/60 \times 1760/a = 88rx/3a$ = number of poles passed in x minutes, or while the train is running $rx/60$ miles.

$\therefore 88rx/3a = r$; whence $x = 3a/88$.

The number of minutes depends upon the distance the poles are apart irrespective of the rate of the train.

Also solved by O. W. ANTHONY, P. S. BERG, A. H. HOLMES, C. D. SCHMITT, H. C. WILKES, B. F. YANNEY, and G. B. M. ZERR.

61. Proposed by COOPER D. SCHMITT, M. A., Professor of Mathematics, University of Tennessee, Knoxville, Tennessee.

Demonstrate the identity $2^{2n+1} \frac{d^n}{dx^n} \left(x^{n+\frac{1}{2}} \frac{d^{n+1}}{dx^{n+1}} e^{Vx} \right) = e^{Vx}$.

I. Solution by O. W. ANTHONY, M. Sc., Professor of Mathematics in New Windsor College, New Windsor, Maryland.

It may be proved inductively that $\frac{d^n}{dx^n} e^{Vx} = \frac{1}{4x} \frac{d^{n-2}}{dx^{n-2}} e^{Vx} - (n - \frac{3}{2}) \frac{d^{n-1}}{dx^{n-1}} e^{Vx}$.

Change n to $n+3$; then

$$\frac{d^{n+3}}{dx^{n+3}} e^{Vx} = \frac{1}{4x} \frac{d^{n+1}}{dx^{n+1}} e^{Vx} - (n + \frac{3}{2}) \frac{d^{n+2}}{dx^{n+2}} e^{Vx}.$$